

NITROUS OXIDE (Commentary)**How effective is nitrous oxide when used in young children ?**

We owe an unacknowledged debt of gratitude to the French for doing the first really comprehensive study of the use of Entonox[®] (a 50:50 mixture of oxygen and nitrous oxide) to alleviate procedural pain in children. A report on its use on over a thousand occasions (Annequin *et al.*, 2000) was followed a year later by a report on its use to provide pain relief during seven and a half thousand procedures (Gall *et al.*, 2001). Serious problems (cyanosis, bradycardia, apnoea or sustained loss of verbal contact) were only encountered on twenty five occasions, and almost all of these problems occurred in children who had also had benzodiazepine sedation as well as an opiate. Even then the problem resolved within minutes once Entonox use was stopped. Entonox seemed to provide substantial pain relief for a whole range of brief but painful procedures such as suturing, burn dressing, fracture reduction, bone marrow aspiration, lumbar puncture, venepuncture, abscess drainage, biopsy procedures, bronchoscopy, gastroscopy, and cast remodelling. Infiltration with lidocaine was also employed where appropriate for procedures such as suturing. Side effects were infrequent. The commonest was brief euphoria. Four per cent suffered nausea and a few of these vomited; three per cent reported transient dizziness or hallucinations. None of these problems persisted for more than five minutes once Entonox administration was stopped. Ninety-three per cent of those old enough to be questioned said they would willingly accept Entonox analgesia again if offered it. Children old enough to cooperate often found a simple or valved mouthpiece more acceptable than a face mask.

There has been some suggestion that equipment capable of delivering up to 70% nitrous oxide in oxygen might offer even more effective pain relief (Kanagasundaram *et al.*, 2001), but there was little evidence of this in a recent very much larger study involving 762 1–17 year old children (Babl, *et al.*, 2008). If a system capable of delivering a variable amount of nitrous oxide is used a fail-safe device must be employed so that there is no risk of the child receiving less than 30% oxygen, and saturation must be checked with a pulse oximeter. The speed with which nitrous oxide delivers analgesia is a major advantage, as is the speed with which patients recover. Those who first introduced this drug's use in children used a demand valve system similar to that traditionally used by women in labour, but most children under five find this difficult to use. Continuous flow systems with scavenging, using either a mask or a mouth piece, work much better in young children, and this approach is now widely used in children of all ages (Babl, *et al.*, 2008).

Only 0.2% of the children in the French study (132/7511) and none of the 762 children in the recent study from Melbourne, Australia, were less than a year old, but there is no good reason to think that Entonox use is likely be less effective or less safe in children younger than this. The French study showed that a nurse can administer Entonox to children more than a year old with proper training as safely and effectively as a doctor. It would be difficult to mount a study to show that Entonox can control pain in the neonate as effectively as it can in older children (any such assumption will need to be taken 'on trust'), but there is certainly a pressing need to get a study mounted to show whether use is equally safe because, while a brief pulse of IV remifentanyl (q.v.) probably provides the best strategy for managing serious short-lived procedural pain in the ventilated baby, we still know very little about how to minimise pain in a non-ventilated baby. Studies certainly suggest that nitric oxide has advantages over midazolam, or even ketamine, as an adjunct to relieving the distress caused by painful procedures in older children (Luhmann *et al.*, 2001; 2006).

Safe use

Upper airway obstruction is extremely uncommon when sedation is achieved using nitrous oxide, but it should never be offered in a setting where this can not be recognised promptly and overcome without delay. . The American Academy of Pediatrics and the American Academy of Pediatric Dentistry recently issued revised joint guidelines for the monitoring and management of children during and after sedation for diagnostic and therapeutic procedures (Coté and Wilson, 2006). The Scottish Intercollegiate Guideline Group had issued rather similar guidance two years earlier (SIGN Guideline 58) which can be downloaded from

www.sign.ac.uk/pdf/sign58.pdf

References

- Griffin G, Campbell V, Jones R. Nitrous oxide-oxygen sedation for minor surgery. Experience in a pediatric setting. *JAMA* 1981;**245**:2411–30.
- Gamis AS, Knapp JF, Glenski JA. Nitrous oxide analgesia in a pediatric emergency department. *Ann Emerg Med* 1989;**18**:177–80.
- Henderson JM, Spence DG, Komocar L, *et al.* Administration of nitrous oxide to pediatric patients provides analgesia for venous cannulation. *Anesthesiology* 1992;**70**:269–71.
- Vetter TA. A comparison of EMLA cream versus nitrous oxide pediatric venous cannulation. *J Clin Anesth* 1995;**7**:486–90.
- Brislin R, Stayer SW, Schwartz R, *et al.* Analgesia for venepuncture in a pediatric surgery centre. *J Paediatr Child Health* 1995;**31**:542–4.
- Evans JK, Buckley SL, Alexander AH, *et al.* Analgesia for the reduction of fractures in children: a comparison of nitrous oxide with intramuscular sedation. *J Pediatr Orthop* 1995;**15**:73–7.
- Gregory PR, Sullivan JA. Nitrous oxide compared with intravenous regional anaesthesia in pediatric forearm fracture manipulation. *J Pediatr Orthop* 1996;**16**:187–91.

- Litman RS, Berkowitz RJ, Wasrd DS. Levels of consciousness and ventilatory parameters in young children during sedation with oral midazolam and nitrous oxide. *Arch Pediatr Adolesc Med* 1996;**150**:671–5.
- Burton J, Auble T, Fuchs S. Effectiveness of 50% nitrous oxide / 50% oxygen during laceration repair in children. *Acad Emerg Med* 1998;**5**:112–7.
- Vic P, Laguette D, Blondin G, *et al.* Utilisation du mélange équimolaire oxygène-protoxyde d'azote dans un service de pédiatrie générale. *Arch Pédiatr* 1999;**6**:844–8.
- Gall O, Annequin D, Ravault N, *et al.* Relative effectiveness of lignocaine-prilocaine emulsion and nitrous oxide inhalation for routine preoperative laboratory te sting. *Paediatr Anaesth* 1999;**9**:305–10.
- Bruce E, Frank L. Self-administered nitrous oxide (Entonox[®]) for the management for the management of procedural pain. *Paediatr Nurs* 2000;**12**:15–9.
- Annequin D, Carbajal R, Chauvin P, *et al.* Fixed 50% oxide oxygen mixture for painful procedures: a French survey. *Pediatrics* 2000;**105**:e47.
- Vater M, Hessell D. Nitrous oxide and oxygen mixture (Entonox[®]) and acute procedure pain. *Paediatr Perinat Drug Ther* 2000;**4**:35–43.
- Pickup S, Pagdin J. Procedural pain. *Paediatr Nurs* 2001;**12**:33–6.
- Krauss B. Continuous-flow nitrous oxide: searching for the ideal procedural anxiolytic for toddlers. *Ann Emerg Med* 2001;**37**:61–2.
- Luhmann JD, Kennedy RM, Porter FL, *et al.* A randomised clinical trial of continuous flow nitrous oxide and midazolam for sedation of young children during laceration repair. *Ann Emerg Med* 2001;**37**:20–7. [RCT] (See also 61–2.)
- Kanagasubndaram SA, Lanae LJ, Cavalletto BP, *et al.* Efficacy and safety of nitrous oxide in alleviating pain and anxiety during painful procedures. *Arch Dis Child* 2001;**84**:492–5.
- Gall O, Annequin D, Benoit G, *et al.* Adverse events of premixed nitrous oxide and oxygen for procedural sedation in children. *Lancet* 2001;**358**:1514–5.
- Kalach N, Barbier C, El Kohen R, *et al.* Évaluation de la tolérance au mélange protoxyde d'azote/oxygène (Entonox[®]) au cours des soins douloureux aux urgences pédiatriques: à propos de 600 observations. *Arch Pédiatr* 2002;**9**:1213–5.
- Lyratzopoulos G, Blain KM. Inhalation sedation with nitrous oxide as an alternative to dental anaesthesia for children. *J Public Health Med* 2003;**25**:303–12.
- Frampton A, Browne GJ, Lam LT, *et al.* Nurse administered relative analgesia using high concentration nitrous oxide to facilitate minor procedures in children in an emergency department. *Emerg Med J* 2003;**20**:410–3.
- Michaud L. Sedation for diagnostic upper gastrointestinal endoscopy: a survey of the Francophone Pediatric Hepatology, Gastroenterology, and Nutrition Group. *Endoscopy* 2005;**37**:167–70.
- Luhmann JD, Schootman , Luhmann SJ, *et al.* A randomized comparison of nitrous oxide plus hematoma block versus ketamine plus midazolam for emergency department forearm fracture reduction in children. *Pediatrics* 2006;**118**:e1078–86. [RCT]
- Coté CJ, Wilson SW, and the Working Group on Sedation. American Academy of Pediatrics and American Academy of Pediatric Dentistry. Guidelines for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures: an update. *Pediatrics* 2006;**118**:2587–602.
- Zier JL, Drake GJ, McCormack PC, *et al.* Case-series of nurse-administered nitrous oxide for urinary catheterisation in children. *Anesth Analg* 2007;**104**:876–9.
- Babl FE, Oakely E, Seaman C, *et al.* High-concentration nitrous oxide for procedural sedation in children: adverse events and depth of sedation. *Pediatrics* 2008;**121**:e528–32.

Commentary posted September 2005

Last updated April 2008